Draft version April 24, 2013
Preprint typeset using LaTeX style emulateapj v. 5/2/11

arXiv:1304.6086v1 [astro-ph.GA] 22 Apr 2013

EVOLUTION OF SECOND GENERATION STARS IN STELLAR DISKS OF GLOBULAR AND NUCLEAR CLUSTERS: ω CENTAURI AS A TEST CASE

Alessandra Mastrobuono-Battisti[1,2] & Hagai B. Perets[1]

Draft version April 24, 2013

ABSTRACT

Globular clusters (GCs) and many nuclear clusters (NCs) show evidence for hosting multiple generations of stellar populations. Younger stellar populations in NCs appear to reside in disk like structures, including the nuclear cluster in our own Galactic center as well as in M31. Kinematic studies of the anomalous globular cluster ω Centauri, thought to possibly be a former dwarf galaxy (or a galactic nucleus), show evidence for its hosting of a central, kinematically cold disk component. These observations suggest that formation of second (or multiple) generation stars may occur in flattened disk like structures. Here we use detailed N-body simulations to explore the possible evolution of such stellar disks, embedded in globular clusters. We follow the long term evolution of a disk like structure similar to that observed in ω Centauri and study its properties. we find that a stellar-disk like origin for second generation stellar populations can leave behind kinematic signatures in properties of the clusters, including an anisotropic distribution, and lower velocity dispersions, and we discuss their implications in constraining the origin of second generations stars and their dynamical age.

Subject headings: methods: numerical, Galaxy: globular clusters: individual (ω Centauri), Galaxy: disk, galaxies: nuclei

1. INTRODUCTION

Nuclear star clusters (NCs), stellar systems similar to globular clusters (GCs), have been found in ~ 75% of the late type spiral galaxies and dwarf ellipticals (Böker et al. 2004; Walcher et al. 2005; Côté, P. et al. 2006; Böker 2010). They contain several distinct stellar populations, characterized by different ages and chemical properties. Among these groups of stars, the youngest ones often form centrally concentrated stellar disks (Van den Bosch 1994; Seth et al. 2006). Such stellar structures are also found in the central pc of the Galaxy and in M31. Their populations of massive young (6 ± 2 Myr and ~ 200 Myr, respectively) stars are arranged in one (or more) distinct stellar disks (Tremaine 1995; Genzel et al. 2000; Levin & Beloborodov 2003; Bender et al. 2005; Lu et al. 2009; Bartko et al. 2010; Genzel et al. 2010).

Similarly, and in contrast with previous paradigm, GCs are not composed of a simple stellar population born at the same time from the same gas. Rather, stars in GCs show significant inhomogeneities in the abundance of light elements and their color magnitude diagrams show several isochrones; thus they consist of multiple stellar generations (MSGs) whose age spreads is estimated to extend up to a few hundred Myrs (see e.g. Gratton et al. 2004, 2012).

The origin of these MSGs is still debated; in one of the models the low velocity gas ejected by primordial asymptotic giant branch (AGB) stars is retained inside the cluster, concentrates at its center and there fragments forming new stars (see Gratton et al. 2012, and reference therein). In this scenario, the initial distribution of second generation stars strongly depends on the initial profile of the collected gas (a somewhat similar model was suggested for the origin of M31 young stellar disk by Chang et al. 2007). For instance, if the gaseous material has some angular momentum, it could produce a flattened structure instead the commonly assumed spherical configuration (Bekki 2011; Mastrobuono-Battisti & Perets 2013), in analogy with what is observed in NCs. This would influence significantly the later dynamical evolution and mixing of older and younger stars, as we show in the following.

One of first clusters where MSGs have been found is ω Centauri (NGC 5139, ω Cen). With mass estimates between 2×10^6 and 5×10^6 $M_\odot$ (see Meylan et al. 1995; van de Ven et al. 2006), ω Cen is the most luminous and massive Galactic GC. It is characterized by a flattened structure (Geyer et al. 1993; van de Ven et al. 2006) and by a wide range of metallicities (e.g. Butler et al. 1978; Norris & Da Costa 1995). Its MSGs are characterized by different spatial distributions (e.g. Freeman & Rodgers 1975; Lee et al. 1999; Hughes & Wallerstein 2000; Hilker & Richtler 2000; Pancino et al. 2000; Ferraro et al. 2002; Bedin et al. 2004; Stanford et al. 2004; Sollima et al. 2005). Dynamically, ω Cen's orbit is peculiar, being tightly bound to the Galactic potential, retrograde and highly eccentric (Dinescu et al. 1999). Moreover, it is one of the favorite candidates for hosting an intermediate-mass black hole, whose mass is estimated to be in the range $13,000 \leq M_\bullet \leq 50,000$ $M_\odot$ (Noyola et al. 2008; van der Marel & Anderson 2010; Miocchi 2010; Jalali et al. 2012). Given these features ω Cen is thought to be the remnant nuclei of a satellite dwarf galaxy that merged with the Milky Way (Freeman 1993; Dinescu et al. 1999; Hughes & Wallerstein 2000; Bekki & Freeman 2003; Böker 2008).

Recently van de Ven et al. (2006) used an extension of the Schwarzschild (1979) method to build axisymmetric dynamical models for ω Cen, using anisotropic velocity distributions. Their best fitting model shows a highly isotropic inner region, which becomes increasingly anisotropic outside of 10 arcmin. The nucleus of ω Cen contains a kinematically cold disk com-

---

[1] Physics Department, Technion - Israel Institute of Technology, Haifa, Israel 32000
[2] INAF - Osservatorio Astrofisico di Arcetri, Largo E. Fermi, 5, 50125 Firenze, Italy

TABLE 1
INITIAL CONDITIONS FOR THE SIMULATION RUNS.

Label	N	$m_{*,1}$ $M_\odot$	$m_{*,2}$ $M_\odot$	M_{disk} $M_\odot$	r_d pc	potential
S1,2	4000	25	0	10^5	4.5	isolated
S3	10000	25	12.5	1.5×10^5	4.5	isolated
S4	8000	12.5	0	10^5	4.5	isolated
S5	4000	25	0	10^5	4.5	tides

NOTE. — S1 and S2 are two different realizations of the same disk; S3 is a simulation of the disk with two mass classes but with the same relaxation time; we run also a case with a disk as massive as in S1,2 but with a bigger number of stars (S4). Finally S5 is a simulation where the cluster orbits a point mass galaxy.

ponent, with a radius between 1 and 3 arcmin, an average flattening of ∼ 0.60, and contributing about ∼ 4% to the total mass of the cluster.

The presence of this disk could be closely related to the existence of MSGs in ω Cen and inside other GCs (Bekki 2011). In particular, second generation stars may form in a disk, and leave behind observable kinematic signatures of their former configuration even after long term dynamical evolution. These signatures may allow us to infer information about their initial distribution, and provide us with constraints on their origin.

In this paper we study the long term evolution of ω Cen-like central disk by means of N-body simulations and explore its structure and potentially observable signatures. This test case is used as a pilot to reveal the possible kinematic signatures of second-generation stellar disks and point to possible clues for the initial presence of a second generation disk in GCs and NCs.

The paper is organized as follows. In Section 2 we describe our choice of the initial conditions, the simulations we run and the code that we use. Then, in Section 3, we show the results obtained. These results and their implications are discussed and summarized in Section 4.

2. THE N-BODY SIMULATION

2.1. *The simulation*

We studied the dynamical evolution of an ω Cen-like central disk structure by means of N-body simulations. These simulations have been done using ϕGrape (Harfst et al. 2006), a high precision, collisional, direct summation code originally configured to run on computer clusters accelerated by GRAPE boards (Makino & Taiji 1998). This code has been recently adapted to run on GPU equipped machines by mean of Sapporo, a CUDA library that calculates force on the GPUs mimicking the behavior of GPAPEs (Gaburov et al. 2009).
The code implements a fourth-order Hermite integrator with a predictor-corrector scheme and hierarchical time stepping. The time-step parameter η and the smoothing length ϵ are the parameters that controls the accuracy of the code. In our simulations, we set $\eta = 0.02$ and $\epsilon = 5 \times 10^{-2}$ pc. The code has been run on the cluster “Tamnun” at the Technion.

2.2. *Initial conditions*

The initial conditions for our simulations were set based on the available observational data. Since the relaxation time of ω Cen is longer than a Hubble time (Harris 2010), mass segregation is not expected to have had significant effects on the system. Its structure is therefore modeled as a single-mass, spherical and isotropic King (1966) cluster. To build



FIG. 1.— Projected isodensity contours plot of the disk on the X-Y plane, at the beginning of the simulation, and going downward, after 2, 5 and 12 Gyr.



FIG. 2.— Cumulative distribution of inclination of stars in the disk at different, equally spaced, times. Going upwards the curves correspond to 0, 1.5, 3, 4.5, 6, 7.5, 9, 10.5 and 12 Gyr.

our model we used the parameters that best fit the cluster observed brightness profile: we set $r_c = 4.6$ pc as the core radius, $W_0 = 5.5$ (Meylan 1987) and 2.5×10^6 $\mathrm{M}_\odot$ as the total mass of the GC (see van de Ven et al. 2006). The N-body representation of the spherical component consists of $N = 100,000$ particles.

In the main simulation we run (S1,2), the disk mass is 10^5 $\mathrm{M}_\odot$, and, as the observed one, has a radial extension of 4.5 pc and a flattening similar to that observed. For its N-body representation we used $N_d = 4000$ single mass particles orbiting counter clockwise in the X-Y plane.

Since we are interested in the evolution of the very central region of ω Cen we neglected the currently observed anisotropy of the cluster.

In addition, at least in the central region, the influence of the tidal field of the Galaxy is negligible as compared to the mutual interaction between particles, thus we simulated the cluster as isolated. Nevertheless, in order to exclude spurious effects caused by these assumptions we run another set of test simulations with different initial parameters for the disk (see Table 1). In particular, we put the cluster on a circular orbit in a point mass potential as massive as the Galaxy within ω Cen's current position. Since the disk evolves in the same way as in the case of the isolated cluster, we can conclude that the presence of the galactic tide doesn't introduce any significant effects. We also run a second realization of the same disk+cluster system and, in order to exclude mass segregation effects, we simulated a disk with two different mass classes (12.5 and 25 $\mathrm{M}_\odot$) with the same relaxation time of the original model. In both cases the evolution of the system doesn't differ significantly from the first case. Finally, we tested the dependence on the chosen number of particles by simulating a disk with $2N$ particles (keeping the same total masses of the disks and cluster, but normalizing for the different relaxation time). Again, we find no significant changes compared with our main simulation. For simplicity, and given the insignificant differences between the various models we tested, we focus here on our main simulation (S1), and show only its results, keeping in mind that they also well represent the results of the other models (see Table 1).



FIG. 3.— The angular momentum of the disk. Upper panel: The evolution of the three components of the angular momentum of the disk particles. The L_x L_y components lay on the initial plan of the disk, while L_z is initially perpendicular to the disk. The disk loses angular momentum that is redistributed to the stars in the cluster. Bottom panel: The dispersion of the three components of the angular momentum.

3. RESULTS

If second generation stars form in a disk, it is possible that their initial distribution leaves observable signatures in the parent dense cluster. Thus, we simulated the evolution of the ω Cen-like central disk for a period of time that, re-scaled to the mass of the particles, is comparable with the typical 12 Gyrs age of Galactic GC. We then analyzed the evolution of the angular momentum of the stars in the disk as well as their orbital parameters in order to find relevant kinematic properties that might be detected in real clusters.

Figure 1 shows the isodensity contours of the disk at the beginning of the simulation and after 2, 5 and 12 Gyr. The disk inflates with time and becomes more spherical but its radial size is only slightly increased. The disk stars remain spatially limited inside ~ 15 pc, while the tidal radius of the



FIG. 4.— Top left panel: the velocity dispersion of the disk at four different evolutionary times, 0, 2, 5 and 12 Gyr; v_z is the velocity component perpendicular to the initial plane of the disk. Top right panel: The final radial (blue thick and black thin solid lines) and the tangential (blue thick and black thin dashed lines) velocity dispersion of the disk and of the cluster as a function of the projected radius. Bottom left panel: The final anisotropy parameter of the disk (solid line) compared with that of the cluster (dashed line), as a function of the projected radius. Bottom right panel: Relaxation time of the central region of the initial simulated system as a function of the radius.

cluster is ~ 80 pc.

The distribution of the average inclination of the disk stars becomes broader with time. After 12 Gyrs it is much more uniform (see the cumulative distribution evolution in Figure 2), i.e. the disk becomes more isotropic with time, in agreement with the results in Figure 1.

Figure 3 illustrates the evolution of the three components of the disk angular momentum (AM, upper panel), showing a consistent decrease with time, and the respective dispersions (bottom panel) that increase with time. At the end of the simulation the average AM is still significant; even after the long term evolution the disk stars still do not show an isotropic distribution. Moreover, as the total AM of the system must be conserved, the cluster structure flattens as it exchanges AM with the disk stars. Interestingly, van den Bergh (2008) found a relation between the morphology of the horizontal branch and the flattening of Galactic GCs. It is possible that our result could be connected to this result, since the shape of the horizontal branch is known to be strongly related to the evolutionary history of the GC and therefore with the presence of MSGs.

While the disk inflates, its velocity dispersion grows with time (see top left panel of Figure 4). Despite that, at the end of the simulation, the velocity dispersion of the disk stars is still smaller than that of the cluster stars. Both the radial and tangential velocity dispersion of the disk and that of the cluster differ, with the former considerably smaller than the latter (see the top right panel of Figure 4). Moreover, the disk remains significantly radially anisotropic in velocity space. Its anisotropy parameter, at the end of the simulation, $\beta = 1 - \sigma_t^2/(2\sigma_r^2)$ is indeed positive at any radius, and has an increasing trend with the distance from the cluster center (see bottom left panel of Figure 4).

Thus, even after 12 Gyrs the flat and the spherical components are not yet mixed and can still be spatially and kinematically distinguished. This suggests that signatures of any putative initial distribution of MSGs formed in a disk-like structure might still be observable today. This is true at least in clusters (or NCs) with long relaxation times as ω Cen, as well as in ω Cen itself, where many stellar generations have been found. Finally, as a result of the evolution of the disk, the whole system becomes tangentially anisotropic with time in the region outside 10 pc, in agreement with what found by van de Ven et al. (2006).

4. DISCUSSION AND SUMMARY

GCs are made of several generation of stars whose origin is still unclear. ω Cen is the first cluster where this feature has been observed. Among its peculiarities, this cluster shows a central disk that could be the remnant of the an initial disk where second generation stars formed. Indeed Bekki (2011) showed that the gas ejected by AGB stars can be retained inside the parent massive star cluster and can fragment into a nested flattened and rotationally supported system of young stars. If so, the flattened structure evolves and may eventually leave signatures of its former presence.

In this work we employed N-body simulations to explore the long-term evolution of such stellar disks embedded in dense stellar cluster, where for the initial conditions we made use of the observed disk like structure in ω Cen .

We verified that such a disk becomes more isotropic with time. However, even after a time comparable with the age of Galactic GCs, the cluster stars and the second generation disk stars are not yet completely mixed. The two populations have different spatial distributions, with the second generation stars still confined only to the central region of the cluster, and do not attain a completely relaxed spherical shape at the end of the simulations. As the second generation disk stars isotropize they exchange angular momentum with the first generation cluster stars, which consequently leads to the slight flattening of the host cluster. It is worth noting that a more massive disk, as e.g. shown to form in simulations (Bekki 2011) is likely to introduce an even larger flattening.

We also find that the kinematic properties of the evolved disk stars change with radial distance. The disk preserves a slight ($\beta \sim 0.05$) radial anisotropy within 4 pc; outside this radius β grows significantly. At the same time the whole system comes to be tangentially anisotropic out of 10 pc, in agreement with observations. In addition, the second generation stars are also characterized by a significantly smaller velocity dispersion than the first generation stars in the central region, which decreases even further with the distance from the cluster center.

In cases where relaxation processes are less efficient, initial conditions are better preserved, as the cluster kinematics still hold their dynamical "memory". This explains why the kinematics of the disk stars show stronger signatures of their original disk structure as a function of increasing radius, at which relaxation times are longer (see bottom right panel of Figure 4 and Figure 21 in van de Ven et al. 2006); indeed, a similar distance dependent effect was suggested to constrain the origin of young stars in the Galactic center (Perets & Gualandris 2010; Madigan et al. 2013). A clear implication/prediction of this issue is that second generation stellar populations in GCs and NCs with longer relaxation times, and/or further out from the center, are likely to show stronger anisotropies in their kinematics. Such regions/clusters would serve as the best test sites for identifying fossilizes signature of a disk/flattened primordial structure.

Finally, the observed signatures of such disks may allow us, to some extent, to determine the dynamical age of the respective stellar population. Since longer evolved disk stars show weaker kinematic signatures, the level of anisotropy provides a dynamical age stamp for this stellar population (see Figures 1, 2 and 4), and might even be translated into the real age of the stellar population, after accounting for the host cluster relaxation time.

Follow up studies exploring larger primordial disk structures than those studied here, beginning with the initial conditions just after second generation star formation (e.g. Bekki 2011; Mastrobuono-Battisti & Perets 2013), will allow us to further explore the generic effects of the long term evolution of second generation stellar disks in a wider range of conditions.

REFERENCES

Bartko H. et al. 2010, ApJ, 708, 834

Bedin L. R., Piotto G., Anderson J., Cassisi S., King I. R., Momany Y., & Carraro G. 2004, ApJ, 605, L125

Bekki, K., & Freeman, K. C. 2003, MNRAS, 346, L11

Bekki, K. 2011, MNRAS, 412, 2241

Bender, R., et al. 2005, ApJ, 631, 280

Böker, T., Sarzi, M., McLaughlin, D. E., van der Marel, R. P., Rix, H.-W., Ho, L. C., & Shields, J. C. 2004, AJ, 127, 105

Böker, T. 2008, AJ, 672, L111

Böker, T. 2010, IAU Symposium, 266, 58

Butler, D., Dickens, R. J., & Epps, E. 1978, ApJ, 225, 148

Chang, P., Murray-Clay, R., Chiang, E., & Quataert, E. 2007, ApJ, 668, 236

Côté, P., et al. 2006, ApJS, 165,57

Dinescu D. I., Girard T. M. & van Altena W. F. 1999, AJ, 117, 1792

Ferraro F. R., Bellazzini M., & Pancino E. 2002, ApJ, 573, L95

Freeman K. C., & Rodgers A. W. 1975, ApJ, 201, L71

Freeman, K. C. 1993, ASP Conf. Ser. 48, The Globular Cluster and Galaxy Connection, ed. G. H. Smith & J. P. Brodie (San Francisco, CA: ASP), 608

Gaburov, E., Harfst, S., & Portegies Zwart, S. 2009, NewAstron. 14 630

Genzel, R., Pichon, C.,Eckart, A., Gerhard, O. E., & Ott, T. 2000, MNRAS 317, 348

Genzel R., Eisenhauer F., & Gillessen S. 2010, Rev. Modern Phys., 82, 3121

Geyer E. H., Nelles B., & Hopp U. 1983, A&A, 125, 359

Gratton, R.G., Sneden, C., & Carretta E. 2004, ARA&A, 42, 385

Gratton, R. G., Carretta, E., & Bragaglia, A. 2012, A&ARv, 20, 50

Harfst, S., Gualandris, A., Merritt, D., Spurzem, R., Portegies Zwart, S., & Berczik, P. 2006, NewAstron. 12, 357

Harris, W. E. 1996 (2010 edition), AJ, 112, 1487

Hilker, M., & Richtler, T. 2000, A&A, 362, 895

Hughes, J., & Wallerstein, G. 2000, AJ, 119, 1225

Jalali, B., Baumgardt, H., Kissler-Patig, M., Gebhardt, K., Noyola, E., Lützgendorf, N., & de Zeeuw, P. T. 2012, A&A, 538, A19

King, I. R. 1966, AJ, 71, 64

Lee Y.-W., Joo J.-M., Sohn Y.-J., Rey S.-C., Lee H.-C., & Walker A. R. 1999, Nature, 402, 55

Levin, Y. & Beloborodov, A. M. 2003, ApJ, 590, L33

Lu, J. R., Ghez, A. M., Hornstein, S. D., et al. 2009, ApJ, 690, 1463

Madigan, A. M. et al. 2013, in prep.

Makino, J., & Taiji, M. 1998, Scientific Simulations with Special-Purpose Computers–the GRAPE Systems, by Junichiro Makino, Makoto Taiji, pp. 248. ISBN 0-471-96946-X. Wiley-VCH, April 1998

Mastrobuono-Battisti, A. & Perets, H. 2013, in prep.

Meylan, G., Mayor, M., Duquennoy, A., & Dubath, P. 1995, A&A, 303, 761

Meylan, G. 1987, A&A, 184, 144

Miocchi P. 2010, A&A, 514, A52

Norris, J. E., & Da Costa, G. S. 1995, ApJ, 447, 680

Noyola, E., Gebhardt, K., & Bergmann, M. 2008, ApJ, 676, 1008

Pancino E., Ferraro F. R., Bellazzini M., Piotto G., & Zoccali M. 2000, ApJ, 534, L83
Perets, H. B., & Gualandris, A. 2010, ApJ, 719, 220
Seth, A. C., Dalcanton, J. J., Hodge, P. W., & Debattista, V. P. 2006, AJ, 132, 2539
Stanford, L. M., Da Costa, G. S., Norris, J. E., & Cannon, R. D. 2004, Mem. S.A.It., 75, 290
Schwarzschild M. 1979, ApJ, 232, 236
Sollima A., Pancino E., Ferraro F. R., Bellazzini M., Straniero O., &Pasquini L. 2005, ApJ, 634, 332
Tremaine, S. 1995, AJ, 110, 628
van den Bergh, S. 2008, AJ, 135, 1731
Van den Bosch, F. C., Ferrarese, L., Jaffe, W., Ford H. C., & O'Connell, R. W. 1994, AJ, 108, 1579
van der Marel, R.P., & Anderson, J. 2010, ApJ, 710, 1063
van de Ven, G., van den Bosch, R. C. E., Verolme, E. K., & de Zeeuw, P. T. 2006, A&A, 445, 513
Walcher, C. J. et al. 2005, ApJ, 618, 237